



ORAL PRESENTATION

Open Access

coBRIM: a phase 3, double-blind, placebo-controlled study of vemurafenib versus vemurafenib + cobimetinib in previously untreated *BRAF*^{V600} mutation-positive patients with unresectable locally advanced or metastatic melanoma (NCT01689519)

Paolo A Ascierto^{1*}, Grant A McArthur², Brigitte Dréno³, James Larkin⁴, Gabriella Liskay⁵, Michele Maio⁶, Mario Mandala⁷, Lev Demidov⁸, Daniil Stroyakovskiy⁹, Luc Thomas¹⁰, Luis de la Cruz-Merino¹¹, Victoria Atkinson¹², Caroline Dutriaux¹³, Claus Garbe¹⁴, Ilsung Chang¹⁵, Stephen P. Hack¹⁵, Antoni Ribas¹⁶

From Melanoma Bridge Meeting 2014
 Naples, Italy. 03-06 December 2014

Background

Combined inhibition of BRAF and MEK is hypothesized to improve clinical outcomes by preventing or delaying onset of resistance observed with BRAF inhibitors alone. This randomized phase 3 study evaluated the combination of the BRAF inhibitor vemurafenib and the MEK inhibitor cobimetinib.

Materials and methods

495 patients were randomly assigned to receive vemurafenib + cobimetinib (60 mg QD, 21 days on/7 days off) or vemurafenib (960 mg BID) + placebo. Eligibility included treatment-naïve *BRAF*^{V600} mutation-positive patients with unresectable locally advanced or metastatic melanoma and adequate performance status and organ function. The primary end point was investigator-assessed progression-free survival (PFS). Safety monitoring included serial cardiac and ophthalmic evaluation and measurement of creatine phosphokinase.

Results

Median PFS was 9.9 months with the combination compared with 6.2 months with the control (HR, 0.51; 95% CI, 0.39-0.68; *P*<0.0001). Objective response rate (ORR)

was 68% in the combination and 45% in the control arm (*P*<0.0001), including complete response in 10% in the combination and 4% of patients in the control group. Subgroup analyses of PFS based on key demographic and tumor characteristics were consistent with PFS in the intent-to-treat population, including those with normal or elevated baseline lactate dehydrogenase (LDH). PFS assessed by independent review was comparable with investigator-assessed PFS. Interim overall survival (OS) data showed an HR of 0.65 (95% CI, 0.42-1.00) but did not cross the prespecified stopping boundary. Compared with vemurafenib alone, the combination was associated with a higher incidence of grade 3 or 4 adverse events (65% vs 59%), with no difference in the rate of adverse events leading to study drug discontinuation (13% vs 12%). Most grade ≥3 events occurred in the first 28 days and resolved quickly. Known MEK inhibitor-related toxicities such as diarrhea, serous retinopathy, elevated creatine phosphokinase, and increased liver transaminase levels were more commonly observed with the combination. The majority was grade 1 or 2, occurred between 1 and 4 months in the treatment course, and resolved quickly. The occurrence of secondary cutaneous neoplasms decreased with the combination (4% vs 18%). Photosensitivity was more common in patients treated with the combination (all grades 32% vs 18%).

¹Istituto Nazionale Tumori Fondazione G. Pascale, Naples, Italy
 Full list of author information is available at the end of the article

Conclusion

Cobimetinib + vemurafenib significantly improved PFS and response rate among patients with $BRAF^{V600}$ -mutant metastatic melanoma, with promising preliminary OS analysis. Most combination-related toxicities are mild or moderate, occur early in treatment, and are manageable by dose modification and supportive care; treatment discontinuation is uncommon.

Clinical trial registration number

NCT01689519.

Authors' details

¹Istituto Nazionale Tumori Fondazione G. Pascale, Naples, Italy. ²Peter MacCallum Cancer Centre, Melbourne, Victoria, Australia. ³Hôtel Dieu Place Alexis Ricordeau, Nantes, France. ⁴The Royal Marsden Hospital, London, UK. ⁵National Institute of Oncology, Budapest, Hungary. ⁶Azienda Ospedaliera Universitaria Senese, Siena, Italy. ⁷Papa Giovanni XXIII Hospital, Bergamo, Italy. ⁸N.N. Blokhin Russian Cancer Research Center, Moscow, Russia. ⁹Moscow City Oncology Hospital 62, Krasnogorsk, Russia. ¹⁰Centre Hospitalier Lyon Sud, Pierre-Bénite, France. ¹¹Hospital Universitario Virgen Macarena, Seville, Spain. ¹²Princess Alexandra Hospital, Woolloongabba, QLD, Australia. ¹³Hopital Saint André, Bordeaux, France. ¹⁴University of Tübingen, Tübingen, Germany. ¹⁵Genentech, Inc, South San Francisco, CA, USA. ¹⁶Jonsson Comprehensive Cancer Center at the University of California, Los Angeles, Los Angeles, CA, USA.

Published: 15 January 2015

doi:10.1186/1479-5876-13-S1-O4

Cite this article as: Ascierto *et al.*: coBRIM: a phase 3, double-blind, placebo-controlled study of vemurafenib versus vemurafenib + cobimetinib in previously untreated $BRAF^{V600}$ mutation-positive patients with unresectable locally advanced or metastatic melanoma (NCT01689519). *Journal of Translational Medicine* 2015 **13**(Suppl 1):O4.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

